

# Matrix string interactions

Vipul Periwal and Øyvind Taffjord

*Department of Physics, Princeton University, Princeton, New Jersey 08544*

String configurations have been identified in compactified Matrix theory at vanishing string coupling. We show qualitatively how the interactions of these strings are determined by the Yang-Mills gauge field on the worldsheet. At finite string coupling, this suggests the underlying dynamics is not well-approximated as a theory of strings. This may explain why string perturbation theory diverges badly, while Matrix string perturbation theory presumably has a perturbative expansion with properties similar to the strong coupling expansion of 2d Yang-Mills theory.

String perturbation theory diverges. This divergence has been linked by Shenker [1] to the unusual strength of non-perturbative effects in string theory. Witten [2] has pointed out that the existence of Ramond-Ramond charged states may lead to precisely such non-perturbative effects.

Following the conjecture of Banks, Fischler, Shenker and Susskind [3], and using the result of Taylor [4], string configurations have been identified in 2d maximally supersymmetric Yang-Mills theory [5–7]. For another approach to string configurations in Yang-Mills theory, see [8]. What is remarkable about these models [5–7] is that weak string coupling is related to strong Yang-Mills coupling, most explicitly stated in [7]. Strong-coupling expansions typically have a finite radius of convergence [9], so it is of interest to identify how string perturbation theory differs from Matrix string perturbation theory. To address this issue, it is necessary to identify how the interactions of Matrix strings arise from the underlying Yang-Mills theory.

For concreteness, we focus on the identification of string configurations in the limit of vanishing string coupling proposed by Dijkgraaf, Verlinde and Verlinde [7]. An equivalent description of these configurations was given in [5,6]. Recall that, according to Taylor [4], Matrix theory [3] compactified on a circle is described by a 2d sYM theory on  $\mathbf{R} \times$  the dual circle. The action is just the dimensionally reduced action obtained from 10d sYM theory. The 8 components of the 10d gauge potential become scalar matter fields on the 2d worldsheet, corresponding to the the transverse space coordinates of the string in lightcone gauge. The action of interest is

$$S = \frac{1}{2\pi\alpha'} \int \text{Tr} \left( D_\mu X D^\mu X + \Theta^T \Gamma^\mu D_\mu \Theta + g_s^2 F_{\mu\nu} F^{\mu\nu} + \frac{1}{g_s^2} [X^i, X^j]^2 + \frac{1}{g_s} \Theta^T \Gamma^i [X_i, \Theta] \right),$$

where  $i = 1, \dots, 8$ ,  $\mu = 0, 1$ , are worldsheet indices, and  $\Theta$  is a Majorana-Weyl spinor in 10d. The worldsheet is taken to be cylindrical with  $\sigma \in [0, 2\pi]$ . This is to be considered for large  $N$ , where  $U(N)$  is the gauge group, following [3].

It was argued in [5–7] that at  $g_s = 0$  the equations of motion imply that the  $X$  matrices mutually commute, with  $[\Theta, X] = 0$ , and hence may be diagonalized simultaneously. The limit  $g_s = 0$  should correspond to an infrared fixed point, and hence to a superconformal field theory. As one goes from  $\sigma = 0$  to  $\sigma = 2\pi$ , the eigenvalues may have nontrivial monodromy, in the sense that the eigenvalues only label orbits of  $U(N)$  up to permutations, so if  $\lambda^i$  stands for the diagonalized  $X^i$  matrices,

$$\lambda(\sigma = 0) = P \lambda(\sigma = 2\pi) P^{-1},$$

for some permutation matrix  $P$  in the defining representation of  $S_N$ . Ref. [5–7] suggested that string configurations should correspond to cycles in this permutation.

The interaction of strings in the model was explained in [7] as arising from the restoration of a non-Abelian  $U(2)$  subgroup when two eigenvalues coincide. Surprisingly, the description of this interaction given in [7], in terms of a twist field in the superconformal field theory that describes the IR fixed point at  $g_s = 0$ , makes *no* reference to the Yang-Mills Lagrangian. Of course, if there is no contact with the Yang-Mills Lagrangian, the Matrix approach would be a particularly obtuse way of thinking about light-cone string perturbation theory (which is not an enlightening approach to string theory in the first instance). In particular, it is unclear why configurations of a few long strings should dominate the dynamics from the given description of string interactions. The only non-perturbative content in the model is the Yang-Mills action, and one would like to *derive* the conjectured string interaction from this action. This should also dynamically determine which configurations of strings actually dominate the dynamics.

We show in this note how the Yang-Mills gauge field on the worldsheet determines the dominant transitions between different string configurations in this model. In doing so, we find that the eigenvalue description is really only suited to  $g_s = 0$ . Away from  $g_s = 0$ , the physics is much clearer in terms of the full matrices. We believe this is the reason why string perturbation theory diverges badly—it is simply that the description in terms of just string configurations, *i.e.* the eigenvalues in [5–7], is valid only at  $g_s = 0$ !

We are interested in the sYM theory at strong Yang-Mills coupling. Expanding  $S$  about an ultralocal theory, we observe that, due to supersymmetric cancellations, the first quantum corrections to the action appear at order  $g_s^2$

for configurations with commuting  $X$  matrices. It is therefore meaningful to consider the effects of classical terms of order  $g_s^0$ . These terms imply that if the  $X$  matrices commute,  $A$  is that connection such that the  $X$  matrices are covariantly constant. Since the  $X$  matrices are sections of a twisted bundle, with monodromy  $P$ , the holonomy of the gauge connection must also be  $P$  for minimizing the  $D_\mu X D^\mu X$  term in the action. While we expressed this in the continuum formulation, it is even simpler to see this from the form of the lattice covariant derivative,  $D_\mu X(x) \equiv U_{x,\mu} X(x + e_\mu) U_{x,\mu}^{-1} - X(x)$ .

Since the 2d sYM theory is strongly-coupled, an appropriate starting point for calculations is the lattice theory. Continuum weak-coupling engineering dimensions of operators are not meaningful in such a lattice theory, so we will not be able to derive the dimension of the operator in the conformal theory ( $g_s = 0$ ) that generates interactions. It is possible, however, to ask qualitative questions about the nature of interactions implied by the Yang-Mills action.

We deduce the *approximate* form of the dominant transitions from the remaining non-vanishing term in  $S$ : The transition from a configuration described by a monodromy  $P$  to one with monodromy  $P'$  is weighted approximately (in Euclidean lattice gauge theory terminology) by

$$\exp \left( \frac{g_s^2}{2\pi\alpha'} \text{Tr}(P' P^{-1} + P P'^{-1} - 2) \right)$$

in  $A^0 = 0$  ‘gauge’. Thus the curvature of the gauge field induced by a transition from one monodromy to another is the determining factor in the relative strengths of different interactions. When the monodromy matrices are permutation matrices, it is easy to see that the least possible curvature is induced by a transition in which two cycles coalesce to form a longer cycle, or vice versa. Comparing this to the interactions described in [7], we see that the sYM action provides a qualitative justification for the assumed form of string interactions.

This is, of course, a gross oversimplification, and should not be taken to mean that  $F^2$  is the continuum operator that generates interactions in the strong-coupling conformal field theory. Indeed, the spectrum of scaling operators at  $g_s = 0$  has little to do with the different terms in  $S$ . We have only deduced the qualitative form of the interactions of string configurations from the sYM action. The other terms that appear at the same order ( $g_s^2$ ) are (1) quantum corrections to the ultralocal theory, and (2) quantum corrections from fluctuations of the terms that determined the holonomy of the connection. To determine the precise operator form of interactions at  $g_s \neq 0$ , one must include these quantum effects. In particular, the transition from one string configuration to another is affected by off-diagonal matrix elements of  $X$ , and by the fluctuations of the gauge field on the world-sheet. The dimension 3 string splitting operator [7] that appears exactly at  $g_s = 0$  is a result of all these dynamical effects. For  $g_s > 0$ , one is dealing with the full matrix structure of the theory, and the stringy nature is no longer obvious.

For a more systematic exploration of the strong-coupling limit of the lattice gauge theory, it would be appropriate to rescale the fields in the action. Written as a dimensional reduction of 10d sYM theory, we would expect just to see  $g_s^2$  in front of the entire action. By rescaling  $X, \Theta$  one can write the action as  $S$  above, or as

$$S' \equiv \frac{1}{2\pi\alpha'} \int \text{Tr} \left( g_s D_\mu X D^\mu X + \sqrt{g_s} \Theta^T \Gamma^\mu D_\mu \Theta + g_s^2 F_{\mu\nu} F^{\mu\nu} + [X^i, X^j]^2 + \Theta^T \Gamma^i [X_i, \Theta] \right),$$

a form suitable for an expansion in  $g_s$  since all terms involving derivatives have positive powers of  $g_s$ . This is equivalent to a strong-coupling expansion in the Yang-Mills coupling.

The lack of a clear stringy interpretation for Matrix string interactions is a good thing, since if the stringy description were to be valid at  $g_s \neq 0$ , we would be left with firm evidence *against* Matrix theory, given the different perturbative behaviours of strong coupling Yang-Mills theory, describing Matrix strings, and garden-variety string theory. If one could surmount the problem of formulating supersymmetric lattice gauge theories [10], one would have a concrete tool to understand Matrix string theory, and see how its behaviour differs from string theory.

Conversations with Wati Taylor and Larus Thorlacius are gratefully acknowledged. This work was supported in part by NSF grant PHY96-00258.

- 
- [1] S.H. Shenker, in *Random surfaces and quantum gravity*, (O. Alvarez, E. Marinari and P. Windey, eds.), Plenum Press (New York, 1991)
  - [2] E. Witten, Nucl. Phys. B443, 85 (1995)
  - [3] T. Banks, W. Fischler, S. Shenker and L. Susskind, Phys. Rev. D55, 5112 (1997)
  - [4] W. Taylor, Phys. Lett. B394, 283 (1997)

- [5] L. Motl, hep-th/9701025
- [6] T. Banks and N. Seiberg, Nucl. Phys. B497, 41 (1997)
- [7] R. Dijkgraaf, E. Verlinde and H. Verlinde, Nucl. Phys. B500, 43 (1997)
- [8] N. Ishibashi, H. Kawai, Y. Kitazawa, A. Tsuchiya, hep-th/9612115
- [9] J.-M. Drouffe and J.-B. Zuber, Phys. Rep. 102, 1 (1983)
- [10] A particularly clear discussion is given in M.F.L. Golterman and D.N. Petcher, Nucl. Phys. B319, 307 (1989); for a recent survey, see I. Montvay, hep-lat/9709080